

Ivor M. Hughes

Barrister & Solicitor

*Patent & Trade Mark Agents
Canada, United States*

*Barrister & Solicitor
Ivor M. Hughes*

*Patent Agents
Neil H. Hughes, P.Eng.
Marcelo K. Sarkis, P.Eng.*

*Counsel
Alfred Schorr*

Our Ref.: PT-1817

December 16, 1999

BY COURIER

The Commissioner of Patents,
UNITED STATES PATENT OFFICE,
2011 South Clark Place,
Crystal Plaza 2, Room 1B03
Arlington, Virginia. 22202
U.S.A.

Dear Sir:

**Re: United States Patent Application
of Kenneth Stephen Albert and Paul José Maes
for CHRONOTHERAPEUTIC FORMULATIONS OF DILTIAZEM AND THE
ADMINISTRATION THEREOF**

Enclosed herewith please find the following documentation for filing with the Commissioner:

- (a) Declaration, Power of Attorney and Petition,
- (b) Abstract in duplicate,
- (c) Disclosure,
- (d) Claims in duplicate,
- (e) Informal Drawings,
- (f) Data Entry Form.


Enclosed along with this material please find a cheque in the amount of **\$1,956.00 U.S. dollars** which includes \$760.00 for the base filing fee of a large entity, \$936.00 for (52) claims over and above the twenty claims allowed per application and \$260.00 for multiple dependent claims. If there is any deficiency or surplusage of the fees enclosed for filing this Application, please obtain any such deficiency or credit the surplusage to Deposit Account 08-3255 and advise Applicants' Agent.

Also enclosed herewith is a stamped, self-addressed verification card which we request that you kindly acknowledge and return to this office at the earliest



opportunity. We thank the Commissioner for his cooperation in this regard and look forward to receiving filing data in this matter.

Respectfully submitted,



Ivor M. Hughes
Agent for Applicant

IMH*kdk
Enclosures

654721-8889460

Initial Information Data Sheet

Inventor Information

Inventor One Given Name :: Kenneth S.
Family Name :: Albert
Postal Address Line One :: 87 Greenwood Lane
Postal Address Line Two ::
City :: Mt. Kisco
State/Province :: NY
Postal or Zip Code :: 10549
Citizenship :: United States
Inventor Two Given Name :: Paul J.
Family Name :: Maes
Postal Address Line One :: 1242 King's College Drive
Postal Address Line Two ::
City :: Oakville
State/Province :: Ontario
Postal or Zip Code :: L6M 2T9
Citizenship :: Belgian

Correspondence Information

Name Line One :: Ivor M. Hughes
Address Line One :: 175 Commerce Valley Drive West
Address Line Two :: Suite 200
City :: Thornhill
State/Province :: Ontario
Country :: Canada
Postal or Zip Code :: L3T 7P6
Telephone :: (905) 771-6414
Fax :: (905) 771-6420

Application Information

Title Line One :: Chronotherapeutic Formulations of
Title Line Two :: Diltiazem and the Administration
Title Line Three :: Thereof
Total Drawing Sheets :: 8
Formal Drawings :: N
Application Type :: Utility

Representative Information

Registration Number One :: 27,759
Registration Number Two :: 33,636
Registration Number Three :: 37,015

Prior Foreign Applications

Foreign Application One ::
Filing Date :: December 10, 1999
Country :: Canada
Priority Claimed :: Y

Separator Sheet



AAA

Application Text

Place Papers behind this sheet in the following order:

1. Preliminary Amendment
2. Specification (English Language Only)
3. Claims
4. Abstract

TITLE OF INVENTION

CHRONOTHERAPEUTIC FORMULATIONS OF DILTIAZEM AND THE
ADMINISTRATION THEREOF

FIELD OF INVENTION

5 This invention relates to once daily preparations comprising Diltiazem and
pharmaceutically acceptable salts thereof, such as the hydrochloride salt, suitable
for evening administration to patients suffering hypertension and/or angina.
This invention also relates to a method for evening administration of such once
daily preparations to patients for the treatment of the patients' hypertension
10 and/or angina.

BACKGROUND OF THE INVENTION

Diltiazem, a benzothiazepine, is an orally active calcium channel blocker
(calcium-antagonist) with relatively high selectivity for vascular smooth muscle
that is effective in the treatment of hypertension and angina pectoris. Today,
15 persons having these conditions take prescribed once daily preparations of
Diltiazem generally to maintain constant levels of the drug in the body over a 24-
hour period. Until recently the timing of the taking of the medicine wasn't
considered an important consideration by the medical community. Doctors
generally did not take into account the natural circadian variation in the body's
20 physiological functions. Researchers have now found that the timing of the
taking of a medicine can affect the way the human body responds to the
medicine. The science of treating the human body taking into account the
natural circadian variation is Chronotherapeutics. Chronotherapeutics relies on
the practice of delivering the correct amount of medication to the correct site of
25 action at the most appropriate time period for the particular disease or condition.

In man, blood pressure does not remain constant during day and night. Early in the morning blood pressure begins to rise from the low levels reached during sleep. Increases in blood pressure are accompanied by increases in heart rate caused by the chemicals generated by the body and delivered into the blood stream. Epidemiological studies have indicated that the greatest incidence of heart problems such as stroke, heart attack, myocardial ischemia and sudden cardiac death occur during the early morning waking hours when the blood pressure is rising in response to the natural circadian rhythm. After normally rising in the morning, blood pressure remains elevated during the day until generally early evening when it starts to fall to its lowest level during sleep.

In one study, evening medication with Diltiazem for treatment of hypertension for effect the next morning has been stated to be more efficacious than other dosage schedules. *Administration Time - Dependent Effects of Diltiazem on The 24-Hour Blood Pressure Profile of Essential Hypertension Patients*, Isao Kohno et al. (Chronobiology International 14(1), 71-84, (1997).) In the report of the study, Herbesser RTM (200mg) was identified as the Diltiazem preparation. Herbesser RTM is a Diltiazem formulation comprising a mixture of immediate release diltiazem - containing microspheres and sustained release diltiazem - containing coated microspheres. According to the report, following a single dose (200 mg) administration, the time of peak plasma diltiazem concentration occurred at 12.5 hours after administration. The peak plasma diltiazem concentration C_{max} in the persons studied was 107mg/ml. Following multiple dosages of 200 mg Diltiazem given over 7 days, the time of peak plasma diltiazem concentration (C_{max}) was at 10 hours after administration. C_{max} was 154 mg/ml.

However a careful review of the report shows inconsistencies which cannot support the authors' conclusions. Particularly at page 80, the best results shown in the graph are with respect to morning treatment with this formulation. Moreover at page 82, the authors themselves acknowledge the study cannot lead to reliable conclusions "because the number of patients was too small". Further, an immediate release portion of the dosage in the order of 15% is not desirable for evening administration. When the blood pressure is naturally at its lowest, not only is there no need for further reduction at that time, but such reduction can harm the patient. Particularly, if the blood pressure is reduced below a minimum the patient is put at a greater risk for cardiovascular accidents including stroke. Further, the 15% immediate release diltiazem is no longer available when needed.

In *A comparative study of the steady-state pharmacokinetics of immediate-release and controlled-release diltiazem tablets*, O. R. Leeuwenkamp et al., Eur. J. Clin. Pharmacol (1994) 46:243-247, controlled release properties and relative systemic availabilities of two dosages of the same controlled release diltiazem tablet formulation were studied by comparing them as steady state with those of an immediate release formulation. In the testing, the diltiazem plasma concentration increased slowly from about 6 hours after the evening dose of both CR tablets (Diltiazem CR 90mg and Diltiazem CR 120 mg) resulting in relatively high plasma concentrations in the early morning hours. The clinicians concluded that twice-daily treatment with diltiazem CR tablets can replace thrice-daily treatment with a conventional diltiazem IR tablet. According to the clinicians "The early morning rise of the diltiazem plasma concentration, which might lead

to a lower incidence of ischemic events, may be an important clinical advantage of both CR tablets."

On April 22, 1998, Searle Canada announced that its Chronovera (R) (controlled onset extended-release verapamil) a high blood pressure medication was now
5 available in Canada. Chronovera (R) was, according to Searle Canada, specifically designed to work with the body's natural circadian variations and was designed to be taken once-a-day just before bedtime. Chronovera provided 24-hour blood pressure control but was designed to deliver peak concentrations of verapamil in the morning when the blood pressure, heart rate and incidence of
10 cardiovascular events were highest. According to Searle Canada, simply changing the time you take the drug your physician has prescribed will not provide the same safety and effectiveness that is designed specially for chronotherapy using verapamil. According to Searle Canada, its Chronovera (R) is unlike traditional medications including extended-release (XL) and sustained-
15 release (SR) formulations which are usually prescribed in doses that maintain relatively constant levels of the drug in the body over a 24-hour period or attempt to maintain relatively constant levels of the drug in the body over a 24-hour period. According to Searle Canada, the prior formulations do not take into account the natural circadian variations in the body's physiological functions.

20

Sustained-release, once-daily diltiazem formulations have been taught which may be considered the traditional medication (according to Searle Canada). They appear not to give the benefits meant to be achieved by chronotherapy.

For example, in *Pharmacokinetic Properties and Antihypertensive Efficacy of Once-*
25 *Daily Diltiazem*, J. G. Kelly et al., *Journal of Cardio-Vascular Pharmacology*,

17:6:957-963, (1991), the controlled-release formulation of diltiazem released a proportion of the diltiazem relatively rapidly with the remainder released over a period extending to 24-hours. During *in vitro* dissolution testing 15% of the diltiazem in the dosage form was released in the first two hours, 54% was released in the first six hours, 89% in the first 13 hours and all of the remainder was released between 13 and 24 hours after administration. The diltiazem capsules contained either 120 mg or 240 mg of diltiazem. It should be noted that no difference is shown between the placebo and dosages in the article at wake-up (between 5:00 a.m. and 8:00 a.m.).

U.S. Patent 4,960,596 discloses slow release 12 hour diltiazem formulations whose dissolution, when measured in accordance with United States Pharmacopoeia 21, purports to be within broad limits (between 5% and 35% after one hour, between 15% and 40% after two hours, between 20% and 50% after three hours, between 30% and 75% after four hours, between 40% and 80% after six hours and between 55% and 95% after eight hours). The examples in the patent, however, provide more specific range limitations specifying range limitations for the formulations exemplified such as at column 4, lines 8-10 and column 5, lines 60-62. In the first series of examples the release into aqueous medium was measured using the method of USP No. 21 of 10%-20% after one hour, 30%-35% after four hours and 60%-75% after eight hours. In the later examples, the release into aqueous medium was measured using the method of USP No. 21 at 15%-35% after one hour, 55%-75% after four hours, 75%-95% after eight hours. These formulations were, however, twice a day (b.i.d.) formulations.

A series of patents have issued to Elan Corporation p.l.c. involving controlled absorption diltiazem pellet formulations for oral administration in which each

pellet has a core comprising diltiazem or a pharmaceutically acceptable salt thereof in association with a specified organic acid covered by an outer membrane which permits release of diltiazem from aqueous medium in accordance with U.S. Pharmacopoeia XX (Paddle Method) in buffered media at
5 pH 1.5, pH 4.0 and pH 7.0. These are U.S. Patent 4,721,619; 4,891,230; 4,894,240; 4,917,899; 5,002,776; 5,219,621; 5,336,504; 5,364,620 and 5,616,345.

In U.S. Patent 4,721,619, dissolution rates of the pellets of examples are found at column 4, lines 41-49 and column 5, lines 5-12. The formulations, however are for 12 hour. The formulations of U.S. Patents 4,891,230 are also for
10 administration every 12 hours.

U.S. Patent 4,894,240 purports to provide formulations for once-daily administration and specifies a general dissolution pattern at column 2, lines 43-52 and a more restricted dissolution pattern at column 3, lines 3-12. The dissolution rates are determined according to U.S. Pharmacopoeia XXI in 0.05M KCl at pH
15 7.0 and at 100 r.p.m. The examples of the patent, however, provide a more limited dissolution pattern under U.S. Pharmacopoeia XXI (Paddle Method) at column 7, lines 30-34 and 47-51, at column 8, lines 16-20, 32-36 and 49-53 and at column 8, line 66 - column 9, line 5. Similar examples are provided at columns 9, 10, 11 and 12. Nothing is taught with respect to formulations suitable as
20 chronotherapeutics.

U.S. Patents 4,917,899, 5,364,620 and 5,616,345 are to the same effect. So are the remaining Elan patents. Nothing in these patents teach formulations suitable as chronotherapeutics.

U.S. Patent 5,529,790 purports to teach a delayed sustained-release
25 pharmaceutical preparation in which a water-soluble drug core is surrounded by

a hydratable diffusion barrier which delays drug release for about two to ten hours. While diltiazem hydrochloride dissolution patterns were provided in accordance with the U.S.P. basket dissolution method specified, no Cmax or the timing of the maximum blood levels is provided. The dissolution rates of the active are not appropriate for a suitable chronotherapeutic (see also U.S. Patents 5,376,384 and 5,478,573).

U.S. Patent 5,288,505 and 5,529,791 relate to extended-release galenical formulations of diltiazem or pharmaceutically acceptable salts thereof which comprise beads in which the active ingredient is in association with a wetting agent and which beads are coated by a microporous membrane. The Cmax of some formulations given in the patents provide for a Cmax after about 8-12 hours. Where the dosing of the formulations of the patents yields maximum diltiazem blood plasma levels (Cmax) of about 145 ng/ml, the Cmax is at about or less than 8 hours.

The applicants are also aware of a formulation marketed under the trade mark Tiazac™ a diltiazem HCl 24-hour sustained-release formulation based on teachings of U.S. Patent 5,529,791 and 5,288,505.

Following chronic administration of Tiazac (240 mg once daily), the average peak plasma Diltiazem concentration (Cmax) is 183 ng/ml (multiple dosage) which occurred after about 7 hours past dose administration. Tiazac™ provides a bioavailability of approximately 59% of the total Diltiazem in the first 12 hours and 41% in the second 12 hours (after 12 hours, 59%; after 16 hours 77% and after 20 hours 90%).

In an article entitled *Effect of Morning Versus Evening Dosing of Diltiazem on Myocardial Ischemia Detected by Ambulatory Electrocardiographic Monitoring in*

Chronic Stable Angio Pectoris, PRA KASH, C. Deedwanian et al., The American Journal of Cardiology, Vol. 80, Aug. 15, 1997, p. 421-425, the authors compare a.m. and p.m. dosing without using an appropriate dosage form for p.m. The Tmax is achieved between 2-6 hours at steady state.

- 5 In an article *The Influence of Time Administration on the Pharmacokinetics of a Once A Day Diltiazem Formulation: Morning Against Bedtime*, Jean Thiffault et al., Biopharmaceutics & Drug Disposition, Vol. 17, 107 - 115 (1996), the once-a-day diltiazem formulation given at 2200 hours for seven days gave "significantly higher plasma concentrations of diltiazem in the early morning hours when the
10 incidence of cardiovascular events is higher". The diltiazem dosages comprise 240 mg taken at 10:00 p.m. (22:00 hours) and maximum concentrations (Cmax) were achieved of 120 ng/ml after about six - eight hours of dosing. Unfortunately, the proposed system covers the period from 2:00 a.m. to 8:00 a.m. To be a true chronotherapeutic, the time period covered should be between about
15 6:00 a.m. and noon. Moreover, this formulation when given at night leads to significantly lower bioavailability than if given in the morning.

It is therefore an object of this invention to provide diltiazem preparations suitable for once-a-day administration in the evening for providing effective dosage amounts in the blood of diltiazem in the morning when blood pressure
20 begins to rise from the low levels reached during sleep, so as to be suitable as a chronotherapeutic preparation.

It is a further object of this invention to provide a method of administration of the diltiazem preparations suitable as a chronotherapeutic so as to be effective in the morning at a time when the patient has most need of the diltiazem preparation.

Further and other objects of the invention will be realized by those skilled in the art from the following summary of the invention and detailed description of embodiments thereof.

SUMMARY OF THE INVENTION

5 According to one aspect of the invention, there is provided a controlled-release Galenical preparation (such as a tablet and a capsule) of pharmaceutically acceptable Diltiazem including the pharmaceutically acceptable salts thereof, such as the hydrochloride salt, suitable for evening dosing every 24 hours containing from about 120 mg to about 540 mg or more (as desired) of the form
10 of Diltiazem associated with excipients to provide controlled (sustained) release of the form of Diltiazem for providing a Cmax of Diltiazem in the blood at between about 10 hours and about 15 hours (preferably about 11 - about 13 hours) after administration, the preparation comprising the form of Diltiazem in oral sustained-release dosage form in which the Diltiazem is adapted to be
15 released after administration over a prolonged period of time and the preparation is adapted to release the Diltiazem

(i) into an aqueous medium at the following rates measured using the method of United States Pharmacopoeia No. XXIII (at 100 rpm in 900 ml of water):

- 20 (a) between about 1% and about 15% after about 2 hours, preferably between about 4% and about 8% after 2 hours;
- (b) between about 7% and about 35% after about 4 hours preferably between about 16% and about 21% after 4 hours;
- (c) between about 30% and about 58% after about 8 hours
25 preferably between about 44% and about 52% after 8 hours;

(d) between about 55% and about 80% after about 14 hours preferably between about 69% and about 76% after about 14 hours; and

(e) in excess of about 75% after about 24 hours and preferably more than about 85% after 30 hours.

and/or (ii) into a buffered medium (such as, for example, phosphate buffer (U.S.P.)) having a pH between about 5.5 and about 6.5, preferably about 5.8 at the following rates measured using the method of United States Pharmacopoeia No. XXIII at 100 rpm in 900ml of the buffered medium:

(a) between about 1% and about 25% after about 2 hours, preferably between about 4% and about 15% after 2 hours;

(b) between about 7% and about 45% after about 4 hours preferably between about 16% and about 30% after 4 hours;

(c) between about 30% and about 68% after about 8 hours preferably between about 44% and about 62% after 8 hours;

(d) in excess of about 75% after about 24 hours and preferably more than 80% after 24 hours.

Preferably no initial retard or delay is built into the preparation retarding /delaying release of Diltiazem from the preparation. Preferably the release rate from the preparation of the Diltiazem is less than about 15% of the total active per hour during dissolution. The preparation may be a diffusion controlled preparation such as, for example, a preparation incorporating the use of microgranules found, for example, in capsules and tablets; tablets; and coated tablets.

The preparation may comprise a plurality of microgranules or pellets, each microgranule comprising a central core or bead containing the form of diltiazem coated with a microporous membrane. The microgranules or pellets may be included in a capsule which dissolves when swallowed to release the

5 microgranules or pellets. The preparation may also comprise a tablet in which the microgranules have been compressed to form the tablet. When compressed into tablet form, wax placebo beads (as known by persons skilled in the art) are preferably included to absorb the shock placed on the microgranules (core and membrane) during the tableting process. By doing so, the integrity of the

10 microgranules containing the Diltiazem active remains intact and the release rate from the preparation is not affected. The tablet may also be coated or uncoated. The preparation may also comprise a sustained-release tablet coating from which preparation the Diltiazem is released. In this regard, the sustained release coating may be applied (sprayed onto) to each tablet.

15 Where the preparation comprises microgranules or pellets (for example) in the capsule or tablet (made, for example, by compressing the microgranules (with preferably wax placebo beads)), the central core may comprise Diltiazem or a pharmaceutically acceptable salt thereof associated with a wetting agent. The Diltiazem may be mixed (in whole or in part) with the wetting agent or may not

20 be mixed with the wetting agent. The wetting agent assists to maintain the solubility of the Diltiazem in each microgranule, ensuring that the solubility of the Diltiazem is unaffected by the pH of the gastrointestinal tract or other adverse conditions which each of the microgranules of the preparation will meet in the gastrointestinal tract.

If the Diltiazem and/or pharmaceutically acceptable salt is not mixed with the wetting agent then the microporous membrane should comprise with suitable adjuvants, a water-dispersible or water-soluble polymer (such as HPMC) and a water-, acid- and base-insoluble polymer of a neutral acrylic polymer such as Eudragit NE30D (a neutral copolymer of acrylic acid ethyl ester and acrylic acid methyl ester) which hydrates the microgranule (including core). If the composition comprises a mixture of the Diltiazem and/or pharmaceutically acceptable salt with the wetting agent, the microporous membrane is preferably the same. However, it may also comprise any suitable membrane which gives to the preparation the required dissolution characteristics.

In this regard, the preferred microporous membrane comprises Eudragit NE30D and hydroxypropylmethylcellulose. This membrane will hydrate the core within the microporous membrane which, for example, may contain diltiazem surrounding a neutral pellet of sugar. The Eudragit NE30D in the membrane expands when it encounters gastrointestinal fluid to greater than 365% of its original size (elongation). This expandability of the membrane gives it the ability to hydrate the membrane and core. The mechanism of release is postulated to be that the membrane will swell while the fluids penetrate and hydrate the core and dissolve the diltiazem and wetting agent. This mechanism is, it is thought, driven by the concentration gradient through the membrane (high concentration inside and low concentration outside).

When Eudragit RS and Eudragit RL are combined to form the microporous membrane, the membrane can expand only very little before breakage or fracturing. The reason is that Eudragit RS expands minimally (about 6%) before the membrane material breaks or fractures changing its release mechanism from

the core. Thus, the mechanism of release from this membrane is thought to be by "washing" the diltiazem through pores created when a plasticizer incorporated in the membrane is released in the gastrointestinal fluid. The diltiazem at the outer surface of the core would be washed from the core through the pores of the microporous membrane, then the diltiazem next presenting itself to the fluids after "washing" of the uppermost (outermost) diltiazem, and so on.

Instead of the wetting agent, any other suitable dissolution agent may be used to assist the release of the Diltiazem from the preparation. For example, instead of the preferred surface active (wetting) agent (surfactant), an organic acid (such as adipic acid, ascorbic acid, citric acid, fumaric acid, malic acid, succinic acid, tartaric acid and the like) may be incorporated in the core. In this regard, the presence of the organic acid in the core permits the diltiazem in the core to dissolve when the composition passes into the higher pH regions of the gastrointestinal tract of the intestine at which pH diltiazem is much less soluble.

One of the membranes which may be used (though not preferred) is the combination of Eudragit RS and Eudragit RL disclosed in U.S. Patent 4,721,619. (See column 1, lines 55-68 and column 2, lines 44-68.) The '619 patent also mentions the use of hydroxypropylmethylcellulose as a water-soluble membrane. The mechanism of release in this case is not by hydration of the core but rather by "washing" the Diltiazem through the pores created in the membrane (for example when the plasticizer in the membrane is released in the gastrointestinal fluid).

The Diltiazem may be present in the core in, for example, the hydrochloride salt form, in which event no dissolution agent may be required in the core.

Suitable preparations such as capsules of the microgranules making up the total Diltiazem active present, may comprise, in the core, Diltiazem hydrochloride between about 50% and about 85% (% w/w of the total preparation (for example, about 69% to about 73%)), a wetting agent (such as sucrose stearate) 5 between about 2% and about 25% (% w/w of the total preparation) (for example about 7% to about 8%) together with suitable adjuvants in the core, and in the membrane between about 0.1% and about 2% of the total preparation of water-soluble and /or water-dispersible polymer such as hydroxypropylmethylcellulose (for example about 0.3% to about 0.6%), and 10 between about 5% and about 20% (% w/w of the preparation) of a neutral copolymer of acrylic acid ethyl ester and acrylic acid methyl ester (such as Eudragit NE30D) (for example about 7% to about 11%).

The microgranules may also be compressed into tablets using suitable excipients. The percentages may be as described above. The tablets may be manufactured, 15 as discussed above, using the microgranules with wax placebo beads and compressing the combination into tablets in the presence of, for example, hydrogenated vegetable oil, sodium starch glycolate and silicone dioxide which have been blended with the microgranules and wax placebo beads before tableting. The tablets may then be coated or uncoated.

20 According to another aspect of the invention, a method of treatment of a patient's hypertension and/or angina is provided comprising administration of a preparation of Diltiazem described above, to the patient in the evening for example at about 7:00 - about 11:00 p.m. for effective treatment of the patient's hypertension and/or angina the next morning, for example between about 6:00 25 a.m. and about noon.

Thus a 24-hour diltiazem preparation is provided wherein the C_{max} of diltiazem in the blood is provided from about 10-15 hours after administration of a single dosage to a patient or about 9-15 hours after multiple dosages over a number of days and displays the dissolution pattern described above determined according to USP 23, page 1791 using Apparatus 1. Apparatus 1 is described as consisting of the following:

a covered vessel made of glass or other inert, transparent material¹; a motor; a metallic drive shaft; and a cylindrical basket. The vessel is partially immersed in a suitable water bath of any convenient size that permits holding the temperature inside the vessel at $37 \pm 0.5^\circ$ during the test and keeping the bath fluid in constant, smooth motion. No part of the assembly, including the environment in which the assembly is placed, contributes significant motion, agitation, or vibration beyond that due to the smoothly rotating stirring element. Apparatus that permits observation of the specimen and stirring element during the test is preferable. The vessel is cylindrical, with a hemispherical bottom. It is 160 to 175 mm high, its inside diameter is 98 to 106 mm, and its nominal capacity is 1000 mL. Its sides are flanged at the top. A fitted cover may be used to retard evaporation.² The shaft is positioned so that its axis is not more than 2 mm at any point from the vertical axis of the vessel and rotates smoothly and without significant wobble. A speed-regulating device is used that allows the shaft rotation speed to be selected and maintained at the rate specified in the individual monograph, within $\pm 4\%$.

Shaft and basket components of the stirring element are fabricated of stainless steel, type 316 or equivalent, to the specifications shown in Figure 1. Unless otherwise specified in the individual monograph, use 40-mesh cloth. A basket having a gold coating 0.0001 inch (2.5 μ m) thick may be used. The dosage unit is placed in a dry basket at the beginning of each test. The distance between the inside bottom of the vessel and the basket is maintained at 25 ± 2 mm during the test.

¹ The materials should not sorb, react, or interfere with the specimen being tested.

² If a cover is used, it provides sufficient openings to allow ready insertion of the thermometer and withdrawal of specimens.
(taken from USP 23)

According to another aspect of the invention, where the preparations comprise cores wherein the diltiazem is in association with a wetting agent, the wetting agent may be selected from:

sugars;

saccharose, mannitol, sorbitol;

lecithins;

C₁₂ to C₂₀ fatty acid esters of saccarose, commercialized under the name of sucroesters (Gattefosse, France) or under the name of crodesters (Croda, U.K.) such as sucrose stearate marketed under the trade name of Crodesta; xylose esters or xylites;

polyoxyethylenic glycerrides;

esters of fatty acids and polyoxyethylene (Brijs, Renex and Eumulgines, Henkel, RFA);

sorbitan fatty acid esters (Span, Atlas, U.S.A.);

polyglycides-glycerides and polyglycides-alcohols esters (Gelucires, Gattefosse, France)

Metal salts such as NaCl or sodium lauryl sulphate

The microporous membrane may be of any suitable material or combination of materials known in the art. Where the wetting agent is in association with the diltiazem in the core and not mixed therewith, the microporous membrane should comprise a water-soluble or water dispersible polymer or copolymer such as hydroxypropylmethylcellulose and a water-, acid- and base-insoluble polymer such as a neutral copolymer of acrylic acid ethyl ester and acrylic acid methyl ester such as Eudragit NE30D. This enables the bead to be hydrated by the introduction of intestinal fluids into the bead hydrating the bead and therefore mixing the diltiazem and the wetting agent. The membrane itself, because of the fluids passing through the membrane, will swell. This membrane acts differently from membranes which do not swell. These other non-hydratable or swellable membranes may be made-up, for example, of water-soluble or water-dispersible polymers or copolymers and a water-, acid- and base-insoluble polymer such as Eudragit RS which swell less easily (owing to the reduced content in quaternary ammonium groups) and are only slightly permeable to active ingredients. This membrane is best suited for coating cores of Diltiazem mixed with a wetting agent or organic acid.

Among materials which may be used to make the microporous membranes, may be mentioned particularly polyacrylates and polymethacrylates of the Eudragit type, ethyl celluloses such as Ethocels from Dow U.S.A. and Aquacoat of FMC U.S.A., hydroxypropylmethylcellulose, hydroxyethylcellulose and hydroxypropylcellulose.

Additionally, adjuvants may be put in the formulation as required such as plastifying agents (plasticizer), pigments, fillers, lubricants and anti-foaming agents. For example talc and/or magnesium stearate may be used as a lubricant, dibutyl sebecate as plasticizer, titanium dioxide as a pigment, Tween 80 as an emulsifier and silicone oil as an anti-foaming agent.

The amount of the microporous membrane is adjusted to provide the sustained release characteristics described.

DETAILED DESCRIPTION OF EMBODIMENTS

The invention will now be illustrated with reference to the following examples and with reference to the following Figures:

Figure 1: is a graph illustrating the Diltiazem Concentration (ng/mL) in the blood after a specified period after a single dose of a 300 mg Diltiazem capsule preparation made according to an embodiment of the invention.

Figure 2: is a graph illustrating the Diltiazem Concentration (ng/mL) in the blood over a 24-hour period after giving multiple doses of the same 300 mg Diltiazem capsules referred to with respect to Figure 1 but over a number of days.

Figure 3: is a graph illustrating dissolution profiles generated according to USP 23 using Apparatus 1 (baskets) at 100 r.p.m. in 900 ml of water for capsule preparations made according to embodiments of the invention (120 mg, 180 mg, 240 mg and 300 mg of Diltiazem active).

Figure 4: illustrates the dissolution profile of a 120 mg capsule preparation of Diltiazem HCl in water according to USP 23 (Apparatus 1 - baskets) according to an embodiment of the invention.

Figure 5: illustrates the dissolution profile of a 120 mg capsule preparation of Diltiazem HCl in gastric fluid according to USP 23 (Apparatus 1 - baskets) according to an embodiment of the invention.

Figure 6: illustrates the dissolution profile of a 120 mg capsule preparation of Diltiazem HCl in intestinal fluid according to USP 23 (Apparatus 1 - baskets) according to an embodiment of the invention.

Figure 7: is a graphic comparison of the blood level concentrations of a preparation (240 mg) made according to an embodiment of the invention and Dilacor (240 mg), a 24-hour oral sustained release dosage form of Diltiazem.

Figure 8: is a graphic comparison of the blood level concentrations of a preparation (240 mg) made according to an embodiment of the invention and Tiazac (240 mg), a 24-hour oral sustained-release dosage form of Diltiazem.

Preparations were manufactured according to the percentages and constituents set out below:

15	Component	% W/W
	(1) Diltiazem hydrochloride	69 - 73
	(2) Microcrystalline cellulose (Avicel ph101)	8 - 9.5
	(3) Povidone K30	1 - 2
	(4) Sucrose stearate (crodesta F150)	7 - 8
20	(5) Magnesium stearate NF	0.5 - 2.5
	(6) Talc USP	0.5 - 5.0
	(7) Titanium dioxide (USP)	0.15 - 0.3
	(8) Hydroxypropylmethylcellulose 2910	0.3 - 0.6
	(9) Polysorbate 80 (tween)	0.01 - 0.025
25	(10) Simeticone C emulsion USP (dry of 30%)	0.01 - 0.015

- | | | |
|------|------------------------------|--------|
| (11) | Eudragit NE30 D (dry of 30%) | 7 - 11 |
| (12) | Purified water USP | 0 |

Two Examples of preparations given the above percentages were made as 120 mg and 180 mg strengths of Diltiazem (as the HCl salt) in capsule form.

5 Example 2

Example 3

Strength

Strength

120 mg capsule

180 mg capsule

- | | | | | |
|----|------|-----------------|------|-----------------|
| | (1) | 120.00 | (1) | 180.00 |
| | (2) | 13.63 - 16.18 | (2) | 20.44 - 24.27 |
| 10 | (3) | 1.7 - 3.41 | (3) | 2.56 - 5.11 |
| | (4) | 11.92 - 13.63 | (4) | 17.88 - 20.44 |
| | (5) | 0.852 - 4.26 | (5) | 1.278 - 6.388 |
| | (6) | 0.852 - 8.52 | (6) | 1.278 - 12.78 |
| | (7) | 0.256 - 0.511 | (7) | 0.383 - 0.767 |
| 15 | (8) | 0.511 - 1.02 | (8) | 0.7665 - 1.533 |
| | (9) | 0.0170 - 0.0426 | (9) | 0.0256 - 0.0639 |
| | (10) | 0.017 - 0.0256 | (10) | 0.0255 - 0.383 |
| | (11) | 11.92 - 18.74 | (11) | 17.886 - 28.106 |
| | (12) | 0 | (12) | 0 |

- 20 240 mg and 300 mg strength preparations in capsule form of Diltiazem (as the HCl salt) were also prepared having the same percentages. They provide the release patterns shown in Figure 3. The dissolution profiles of all of the strengths were generated from biobatches of capsules using Apparatus 1 (baskets) at 100 RPM in 900 ml of water in accordance with USP 23.

Less than 20% of the formulation is dissolved after about four hours (for example between about 16%-21%) with less than about 10% dissolved in the first two hours (for example between about 4% - about 8%). Less than about 50% is released after 8 hours (for example between about 44%-52%). Less than about 73% is released after 14 hours (for example 69%-76%). Preferably in excess of about 85% is released after 24 hours.

Specifically, samples of 120 mg capsules of Diltiazem HCl (made according to the embodiment of the invention) had the following dissolution profile:

Percent Dissolved - Time Elapsed

	2h (%)		4h (%)		8h (%)		14h (%)		24h (%)	
	5	8	19	19	49	49	72	72	88	88
	4	5	16	14	32	44	76	69	93	86
	5	6	18	16	50	49	72	73	88	90
	7	6	21	17	54	48	76	72	92	87
	5	8	17	19	51	50	74	74	92	91
	6	7	18	19	52	52	74	75	90	92
Mean (%)	6		18		50		73		90	
RSD	21.3		10.5		5.1		2.7		2.6	

Samples of 180 mg capsules of Diltiazem HCl (made according to an embodiment of the invention) had the following dissolution profiles:

Percent Dissolved - Time Elapsed

Lapsed Time	2h (%)		4h (%)		8h (%)		14h (%)		24h (%)	
	8	7	21	20	52	52	76	73	91	89
	6	7	19	20	52	51	76	73	93	90
	5	6	16	18	48	50	72	72	89	90
	6	7	19	18	52	49	76	72	98	88
	7	7	20	19	51	51	73	74	91	91
	8	7	20	21	51	51	74	73	92	91
Mean (%)	7		19		51		74		91	
RSD	12.8		7.4		2.5		2.1		1.7	

Samples of 240 mg capsules of Diltiazem HCl (made according to an embodiment of the invention) had the following dissolution profiles:

Percent Dissolved - Time Elapsed

	2h (%)	4h (%)	8h (%)	14h (%)	24h (%)
	6 4	19 16	46 48	73 71	86 86
	6 5	18 15	48 45	70 68	85 84
	5 5	18 17	49 49	71 72	86 88
	4 7	16 18	46 48	68 71	83 87
	6 4	18 15	49 50	70 68	84 84
	6 6	18 17	48 48	70 71	85 86
Mean (%)	5	17	48	70	85
RSD	18.5	7.7	2.9	2.3	1.7

5

Samples of Diltiazem HCl capsules 300 mg (made according to an embodiment of the invention) had the following dissolution profile:

Percent Dissolved - Time Elapsed

	2h (%)	4h (%)	8h (%)	14h (%)	24h (%)
	3 4	16 16	46 45	68 67	83 83
	6 5	17 16	49 45	73 67	90 83
	6 5	16 16	46 46	69 68	84 84
	5 5	16 16	46 46	69 69	83 87
	6 4	17 15	46 45	68 68	82 86
	5 5	17 17	46 47	69 70	84 87
Mean (%)	5	16	46	69	85
RSD	13.2	3.8	2.4	2.3	2.8

10

Additionally, the following Dissolution Profiles were obtained for the samples of 120 mg Diltiazem HCl Capsules:

Medium: Water

Hour	Diltiazem HCl Capsules % Dissolved (Average of 12 capsules)	Range [%]	RSD
2	6	4 - 8	17.6
4	18	14 - 21	9.8
8	50	44 - 54	5.1

14	73	69 - 76	2.8
24	90	86 - 93	2.5

Medium: Gastric Fluid

Hour	Diltiazem HCl Capsules % Dissolved (Average of 12 capsules)	Range [%]	RSD
2	5	3 - 6	18.8
4	16	14 - 18	9.0
8	49	47 - 52	3.5
14	73	71 - 75	1.8
24	87	85 - 89	1.5

Medium: Intestinal Fluid

Hour	Diltiazem HCl Capsules % Dissolved (Average of 12 capsules)	Range [%]	RSD
2	5	3 - 7	26.0
4	17	14 - 20	12.0
8	43	40 - 47	6.1
14	64	53 - 69	8.1
24	78	65 - 85	8.1

5

Other Dissolution Profiles were determined of
embodiments of the invention
(Medium - USP Water)

10 Apparatus: USP #1 (baskets) at 100 rpm

Diltiazem 120 mg Capsules

TIME [h]	2	4	8	14	24
vessel 1	5%	19	49	72	88
vessel 2	4	16	52	76	93
vessel 3	5	18	50	72	88
vessel 4	7	21	54	76	92
vessel 5	5	17	51	74	92
vessel 6	6	18	52	74	90
vessel 7	8	19	49	72	88
vessel 8		14	44	69	86
vessel 9		16	49	73	90
vessel 10	6	17	48	72	87
vessel 11	8	19	50	74	91
vessel 12	7	19	52	75	92

MEAN	6%	18	50	73	90
SD	1.3	1.9	2.6	2.0	2.3
RSD	21.3	10.5	5.1	2.7	2.6
RANGE	4 - 8	14 - 21	44 - 54	69 - 76	86 - 93

(Medium - Gastric)

Apparatus: USP #1 (baskets) at 100 rpm

5

Diltiazem 120 mg Capsules

TIME [h]	2	4	8	14	24
vessel 1	3	14	51	74	88
vessel 2	6	17	48	72	85
vessel 3	5	17	49	72	85
vessel 4	5	15	48	72	87
vessel 5	4		47	71	86
vessel 6	6	18	50	72	86
vessel 7		15	49	73	88
vessel 8	4	14	48	71	86
vessel 9	5	17	51	74	88
vessel 10	6	18	52	74	88
vessel 11	6	18		75	89
vessel 12	5	17	50	73	87
MEAN	5	16	49	73	87
SD	0.9	1.5	1.7	1.3	1.3
RSD	19.0	9.0	3.5	1.8	1.5
RANGE	3 - 6	14 - 18	47 - 52	71 - 75	85 - 89

(Medium - Intestinal)

10

Apparatus: USP #1 (baskets) at 100 rpm

Diltiazem 120 mg Capsules

TIME [h]	2	4	8	14	24
vessel 1	7	19	45	67	81
vessel 2	4	14	40	64	79
vessel 3	7	20	47	69	83
vessel 4	5	19	46	68	83
vessel 5		17	41	58	69
vessel 6		17	45	69	83
vessel 7	4	17	40	53	65
vessel 8	5	17	42	65	78
vessel 9	5			58	73
vessel 10	5	17	47	68	85
vessel 11	4	15	44	64	81

vessel 12	4	15	43	64	81
MEAN	5	17	43	64	78
SD	1.2	2.0	2.7	5.2	6.4
RSD	25.9	13.0	6.1	8.1	8.1
RANGE	3 - 7	14 - 20	40 - 47	53 - 69	65 - 85

- Briefly, the dosages in Examples 1 (120 mg) and 2 (180 mg) and the 240 mg and 300 mg dosages were manufactured by mixing the core (bead) ingredients (diltiazem, microcrystalline cellulose, povidone, sucrose stearate) by introducing the components into a planetary mixer and granulating same and mixing with purified water. The plastic mass was then extruded to provide an extrudate. The extrudate was subsequently spheronized to produce diltiazem spheres in admixture with the wetting agent. The spheres (cores) were dried in an oven and sieved to the appropriate size cores or beads.
- 10 The membrane was prepared by mixing the hydroxypropylmethylcellulose, titanium dioxide, talc, magnesium stearate, Polysorbate 80 and Simethacone C emulsion and thereafter combined with the Eudragit NE30D and water. The spheronized cores were coated with the appropriate thickness of membrane by spraying the cores, coating same. Thus the cores (beads) were coated with the coating suspension to produce the microgranules or pellets. The microgranules or pellets were then dried.
- 15

In more detail the process combines Diltiazem Hydrochloride USP, Microcrystalline cellulose NF (Avicel PH 101), Povidone K30 USP and Sucrose Stearate (Crodesta F160) as follows:

- 20 The following were screened through a 1.9 mm screen and added to a mixer bowl:

Diltiazem

Avicel PH 101

Povidone K30.

To remove large agglomerates, the Crodesta 7.98 kg was screened through a 1.0 - 1.2 mm screen and added to the same mixing bowl. The items were then blended in an AMF blender at 50 RPM. 1 kg of the above dry blend was set aside
5 to be used as dusting powder (Diltiazem Dusting Powder). The remainder of the blend was continued to be blended at 50 rpm until adequately granulated. The granulated material was then loaded into the hopper of an extruder (such as EXDCS100 or EXDS 60). The granulation was extruded and without breaking up the extrudate, the extrudate was collected. The extrudate was then spheronized
10 into the cores (beads) of the desired size and were dusted as desired by the Diltiazem Dusting Powder set aside. The beads were then dried by spreading them on trays and drying in an oven set at about 57°C. The Drying Temp. was in the order of 55-60°C for about 12 hours (in the order of 12-16 hours). The dried cores (beads) were sieved to collect those of appropriate size (0.7 - 1.4 mm).
15 A Eudragit NE30D and hydroxypropylmethylcellulose coating suspension, was made. The following:

- Magnesium Stearate NF
- Talc USP
- Titanium Dioxide USP
- 20 Hydroxypropylmethylcellulose 2910 USP (Pharmacoat 606)
- Polysorbate 80 NF (Tween 80)
- Simethicone C Emulsion USP

and pure water were combined within a Silverson Mixer. Water was first mixed with Polysorbate 80 and the Simethicone. The HPMC was then added, then
25 titanium dioxide, then talc and then the Magnesium Stearate. The mixture was

stored for 2 hours. The Eudragit NE30D was screened through a 0.310 mm sieve and added to the mixture.

The beads were then coated with the suspension by using an Aerocoater™ and spraying the beads (which have been preheated to 26°C) with the coating suspension to achieve the desired thickness (about 0.05 mm). The beads were then dried by spreading on trays and drying at 40-45°C for 10 - 12 hours.

Diltiazem HCl 300 mg capsules made according to an embodiment of the invention were tested in a single dose study to determine their bioavailability, their C_{max} and T_{max}, their rate and extent of absorption. Blood sampling for drug content analysis was carried out at 0.0 (predrug) 1, 2, 3, 4, 5, 6, 8, 10, 12, 14, 16, 18, 24, 30, 36, 42 and 48 hours post-drug. Vital sign and 12 - Lead ECG monitoring were conducted at 0 (predrug) 2, 6, 8 and 12 hours post-drug. The following was determined from the plasma study:

Mean Pharmacokinetic Parameters for Plasma Diltiazem
(n = 41)

Parameter	1 x 300 mg Mean (%CV)
AUC (0-t)(ng.hr/mL)	2703.83 (36.26)
AUC (0-inf.)(ng.hr/mL)	2786.95 (36.39)
C _{max} (ng/mL)	146.33 (38.43)
T _{max} (hours)	13.17 (14.79)
t _{1/2} (hours)	6.96 (17.56)
K _{el} (hour ⁻¹)	0.102 (15.983)

Mean Plasma Diltiazem Concentrations (ng/mL)
(n = 41)

SAMPLE TIME (HOURS)	1 x 300 mg
0.00	0.00 ± 0.00
1.00	0.76 ± 2.20
2.00	4.92 ± 3.87
3.00	10.97 ± 5.92
4.00	20.01 ± 10.77

5.00	33.46 ± 18.39
6.00	70.21 ± 37.03
8.00	95.43 ± 41.50
10.00	110.16 ± 47.43
12.00	132.84 ± 52.04
14.00	139.54 ± 55.11
16.00	126.35 ± 50.23
18.00	105.74 ± 40.86
24.00	62.84 ± 24.20
30.00	43.92 ± 16.94
36.00	25.67 ± 11.46
42.00	13.40 ± 7.37
48.00	7.50 ± 4.46

Mean Pharmacokinetic Parameters for Plasma Diltiazem
(n = 36)

Parameter	Geometric Mean Arithmetic Mean (C.V.)
	1 x 300 mg
AUC (0-t hours)(ng.hr/mL)	2682.87 2872.06 (38.44)
AUC (0-x)(ng.hr/mL)	1955.92 2075.00 (35.63)
AUC (0-infinity)(ng.hr/mL)	2847.57 3055.19 (39.05)
C _{max} (ng/mL)	134.96 144.00 (37.17)
T _{max} (hours)**	13.00 (2.92)
t _{1/2} (hours)*	8.69 (22.85)
K _{el} (hour ⁻¹)*	0.084 (22.860)

5 * These are arithmetic means (CV%).

** This is median (± S.D.).

With reference to Figures 1 and 2, it is clear the 300 mg capsule preparation made according to an embodiment of the invention provides the appropriate Diltiazem blood levels at the appropriate time to be suitable for administration as a chronotherapeutic - being given in the evening to provide effective concentrations of Diltiazem the following morning. This suitability is illustrated with reference to Figures 7 and 8. In Figure 7, the 240 mg Diltiazem preparation

made according to the embodiment of the invention provides elevated blood levels that are effective all morning for effective treatment of the patient with Diltiazem. However, the Dilacor formulation (given either in the evening or the following morning) does not protect the patient from 6:00 a.m. - noon, the more dangerous period.

The same is true with Figure 8. Tiazac given in the morning, does not provide the protection. Further, peak plasma concentrations for Tiazac are achieved after about 7 hours after dose administration.

The preparations according to embodiments may also be made as tablets. The tablets may be made as compressed tablets in the desired strengths (for example 120 mg - 540 mg or more Diltiazem) incorporating the microgranules. The tablets may even be scored to permit division into smaller doses.

Tablets may be made as follows using the microgranules or pellets, wax placebo beads and hydrogenated vegetable oil, sodium starch glycolate and silicone dioxide as follows:

The microgranules of Diltiazem may be the following:

Magnesium Stearate

Talc

Titanium Dioxide

Hydroxypropylmethyl-Cellulose 2910

Polysorbate 80

Simethicone Emulsion

Eudragit NE30D

Diltiazem Hydrochloride

Microcrystalline Cellulose

Povidone K30

Sucrose Stearate

Purified Water

The wax placebo beads may be the following:

5 Microcrystalline Wax NF

Pregelatinized Starch

Sodium Starch Glycolate

Titanium Dioxide

Carbon Dioxide

10 The microgranules, wax placebo beads, hydrogenated vegetable oil, sodium starch glycolate and silicone dioxide may be combined and compressed into the desired strengths of tablets, for example 240 mg, 300 mg and 360 mg tablets.

Briefly, to form the microgranules, Diltiazem HCl, Microcrystalline Cellulose, Povidone 30, Sucrose Stearate may be mixed to form a "dry blend". A 1 kg
15 portion of the dry blend may be removed and stored in a separate labeled container as the Dusting Powder, for use in subsequent manufacturing steps (if desired). Following the removal of the Dusting Powder, Purified Water is added to the dry blend and mixed to create a plastic mass. The plastic mass is extruded through a 1.0 mm screen to form a spaghetti like extrudate. This extrudate is
20 then spheronized into beads. During the spheronization process Dusting Powder is added to dry the beads and provide them with a smooth aspect (if required). The addition of Dusting Powder also prevents the newly spheronized beads from sticking together. The spheronized beads are tray dried for 12 - 16 hours and sieved to select beads that are larger than 0.7 mm and smaller than 1.4
25 mm in diameter.

The beads are loaded into a preheated (40 - 45°C) fluid bed Aerocoater. Coating suspension is applied at an amount of 10% by spray coating. The resulting Diltiazem Microgranules (coated beads) are dried for between 10 - 12 hours and the dried coated beads are sieved to select coated beads that are larger than 0.7 mm and smaller than 1.7 mm in diameter.

For the manufacture of the placebo wax beads, Microcrystalline Wax, Pregelatinized Maize Starch, Sodium Starch Glycolate and Titanium Dioxide are mixed in a high shear mixer and heated to 64°C (jacket temperature 70°C). The resulting melt is cooled by the addition of liquid CO₂ to form the solid starters of the pellets. The pellet starters are mixed and the size is increased by the gradual turning of the impeller for a fixed timeperiod (mixing time is directly related to the impeller speed and the time to reach a temperature of $57 \pm 2^{\circ}\text{C}$). The resulting beads are sieved to select beads larger than 0.7 mm and smaller than 1.4 mm in diameter.

For manufacturing the Diltiazem chronotherapeutic tablets, the placebo wax beads and the microgranules of Diltiazem are blended at a ratio of about 2:3 (placebo wax beads: microgranules of Diltiazem) with Hydrogenated Vegetable Oil (lubricant), Sodium Starch Glycolate (disintegrant) and Silicone Dioxide (lubricant) added. The blend is tableted under low pressure (approximately 6 - 8 Sc) to form the compressed Diltiazem Tablets.

In the compressed tablets, the placebo wax beads serve to absorb the shock placed on the microgranules of Diltiazem during the tableting process. By doing so the integrity of the microgranules remains in tact and the release rate of the diltiazem is not affected.

As many changes can be made to the embodiments of the invention without departing from the scope thereof, it is intended that all material contained herein be determined as illustrative of the invention and not in a limiting sense.

THE EMBODIMENTS OF THE INVENTION IN WHICH AN EXCLUSIVE PROPERTY OR PRIVILEGE IS CLAIMED ARE AS FOLLOWS:

1. A controlled-release Galenical preparation of pharmaceutically acceptable Diltiazem including the pharmaceutically acceptable salts thereof, suitable for evening dosing every 24 hours containing from about 120 mg to about 540 mg or more of the form of Diltiazem with excipients to provide controlled (sustained) release of the form of Diltiazem from the preparation for providing a Cmax of Diltiazem in the blood at between about 10 hours and about 15 hours (Tmax) after administration of the preparation, the preparation being in a sustained-release dosage form in which the Diltiazem is adapted to be control released after administration of the preparation over a period of time and being adapted to release the Diltiazem

(i) into an aqueous medium at the following rates measured using the method of United States Pharmacopoeia No. XXIII at 100 rpm in 900 ml of water:

- (a) between about 1% and about 15% after 2 hours;
- (b) between about 7% and about 35% after 4 hours;
- (c) between about 30% and about 58% after 8 hours;
- (d) between about 55% and about 80% after 14 hours; and
- (e) and in excess of about 75% after 24 hours.

and/or (ii) into a buffered medium having a pH between about 5.5 and about 6.5, at the following rates measured using the method of United States Pharmacopoeia No. XXIII at 100 rpm in 900ml of the buffered medium:

- (a) between about 1% and about 25% after about 2 hours;

- (b) between about 7% and about 45% after about 4 hours;
- (c) between about 30% and about 68% after about 8 hours;
- (d) in excess of about 75% after about 24 hours.

2. A controlled-release Galenical preparation of pharmaceutically acceptable Diltiazem including the pharmaceutically acceptable salts thereof, suitable for evening dosing every 24 hours containing from about 120 mg to about 540 mg or more of the form of Diltiazem with excipients to provide controlled (sustained) release of the form of Diltiazem from the preparation for providing a C_{max} of Diltiazem in the blood at between about 10 hours and about 15 hours (T_{max}) after administration, the preparation being in a sustained-release dosage form in which the Diltiazem is adapted to be control released after administration of the preparation over a period of time and being adapted to release the Diltiazem

(i) into an aqueous medium at the following rates measured using the method of United States Pharmacopoeia No. XXIII at 100 rpm in 900 ml of water:

- (a) between about 4% and about 8% after 2 hours;
- (b) between about 16% and about 21% after 4 hours;
- (c) between about 44% and about 52% after 8 hours;
- (d) between about 69% and about 76% after 14 hours; and
- (e) and in excess of about 85% after 24 hours;

and/or (ii) into a buffered medium having a pH about 5.8 at the following rates measured using the method of United States Pharmacopoeia No. XXIII at 100 rpm in 900ml of the buffered medium:

- (a) between about 4% and about 15% after 2 hours;

- (b) between about 16% and about 30% after 4 hours;
- (c) between about 44% and about 62% after 8 hours;
- (d) in excess of about 80% after 24 hours.

3. The preparation of claim 1 or 2 wherein the Cmax of Diltiazem in the blood is obtained between about 11 - about 13 hours after administration of the preparation.
4. The preparation of claim 1, 2 or 3 wherein the Diltiazem is in the form of Diltiazem HCl.
5. The preparation of claim 1, 2, 3 or 4 wherein the preparation is a diffusion controlled preparation.
6. The preparation of claim 1, 2, 3, 4 or 5 wherein the preparation releases the Diltiazem at a rate of less than about 15% of the total amount of active per hour during dissolution.
7. The preparation of claim 1, 2, 3, 4, 5 or 6 in capsule form.
8. The preparation of claim 1, 2, 3, 4, 5 or 6 in tablet form.
9. The preparation of claim 1, 2, 3, 4, 5, 6, 7 or 8 wherein the preparation comprises a plurality of microgranules, each microgranule comprising a central core containing the form of diltiazem coated with a microporous membrane and the central core comprises Diltiazem or pharmaceutically acceptable salt thereof associated with a wetting agent.

10. The preparation of claim 9 wherein the Diltiazem is mixed (in whole or in part) with the wetting agent.
11. The preparation of claim 10 wherein the wetting agent assists to maintain the solubility of the Diltiazem in each bead, ensuring that the solubility of the Diltiazem is unaffected by the pH of the gastrointestinal tract or other adverse conditions which the composition will meet therein.
12. The preparation of claim 9, 10 or 11 wherein the membrane comprises a water-dispersible or water-soluble polymer (such as HPMC) and a water-, acid- and base-insoluble polymer of a neutral acrylic polymer such as Eudragit NE30D (a neutral copolymer of acrylic acid ethyl ester and acrylic acid methyl ester) which hydrates the preparation.
13. The preparation of claim 9 wherein the preparation comprises a mixture of the Diltiazem and/or pharmaceutically acceptable salt with the wetting agent and the membrane comprises a water-dispersible or water-soluble polymer (such as HPMC) and a water-, acid- and base-insoluble polymer of a neutral acrylic polymer such as Eudragit NE30D (a neutral copolymer of acrylic acid ethyl ester and acrylic acid methyl ester) which hydrates the preparation.
14. The preparation of claim 9, 10, 11, 12 or 13 wherein the membrane comprises Eudragit NE30D and hydroxypropylmethylcellulose.

15. The preparation of claim 14 wherein the membrane hydrates the core within a membrane which when put in gastrointestinal fluid causes the membrane to swell while fluid penetrates and hydrates the bead, and dissolves the diltiazem and wetting agent and benefits from a concentration gradient through the membrane (high concentration inside and low concentration outside).

16. The preparation of claim 10 or 11 wherein the Diltiazem is mixed with the wetting agent and the membrane comprises Eudragit RS, Eudragit RL and plasticizer combined to form the membrane thereby providing a mechanism of release from this membrane which "washes" the diltiazem through pores created when the plasticizer incorporated in the membrane, is released in gastrointestinal fluid.

17. The preparation of claim 1, 2, 3, 4, 5, 6, 7 or 8 wherein the preparation comprises a plurality of microgranules comprising a central core containing the form of diltiazem coated with a microporous membrane and the central core comprises Diltiazem or a pharmaceutically acceptable salt thereof associated with any suitable dissolution agent (other than a wetting agent) to assist in the release of the Diltiazem from the preparation.

18. The preparation of claim 17 wherein the dissolution agent is an organic acid such as adipic acid, ascorbic acid, citric acid, fumaric acid, malic acid, succinic acid, tartaric acid and the like which permits the diltiazem to dissolve in gastrointestinal fluids when the microgranules pass into the higher pH regions of the gastrointestinal tract of the intestine at which pH diltiazem is much less soluble.

19. A method of treatment of a patient's hypertension and/or angina comprising the administration of the preparation of Diltiazem of claim 1 to the patient in the evening for effective treatment of the patient's hypertension and/or angina the next morning.

20. A method of treatment of a patient's hypertension and/or angina comprising the administration of the preparation of Diltiazem of claim 2 to the patient in the evening for effective treatment of the patient's hypertension and/or angina the next morning.

21. A method of treatment of a patient's hypertension and/or angina comprising the administration of the preparation of Diltiazem of claim 3 to the patient in the evening for effective treatment of the patient's hypertension and/or angina the next morning.

22. A method of treatment of a patient's hypertension and/or angina comprising the administration of the preparation of Diltiazem of claim 4 to the patient in the evening for effective treatment of the patient's hypertension and/or angina the next morning.

23. A method of treatment of a patient's hypertension and/or angina comprising the administration of the preparation of Diltiazem of claim 5 to the patient in the evening for effective treatment of the patient's hypertension and/or angina the next morning.

24. A method of treatment of a patient's hypertension and/or angina comprising the administration of the preparation of Diltiazem of claim 6 to the patient in the evening for effective treatment of the patient's hypertension and/or angina the next morning.

25. A method of treatment of a patient's hypertension and/or angina comprising the administration of the preparation of Diltiazem of claim 7 to the patient in the evening for effective treatment of the patient's hypertension and/or angina the next morning.

26. A method of treatment of a patient's hypertension and/or angina comprising the administration of the preparation of Diltiazem of claim 8 to the patient in the evening for effective treatment of the patient's hypertension and/or angina the next morning.

27. A method of treatment of a patient's hypertension and/or angina comprising the administration of the preparation of Diltiazem of claim 9 to the patient in the evening for effective treatment of the patient's hypertension and/or angina the next morning.

28. A method of treatment of a patient's hypertension and/or angina comprising the administration of the preparation of Diltiazem of claim 10 to the patient in the evening for effective treatment of the patient's hypertension and/or angina the next morning.

29. A method of treatment of a patient's hypertension and/or angina comprising the administration of the preparation of Diltiazem of claim 11 to the patient in the evening for effective treatment of the patient's hypertension and/or angina the next morning.

30. A method of treatment of a patient's hypertension and/or angina comprising the administration of the preparation of Diltiazem of claim 12 to the patient in the evening for effective treatment of the patient's hypertension and/or angina the next morning.

31. A method of treatment of a patient's hypertension and/or angina comprising the administration of the preparation of Diltiazem of claim 13 to the patient in the evening for effective treatment of the patient's hypertension and/or angina the next morning.

32. A method of treatment of a patient's hypertension and/or angina comprising the administration of the preparation of Diltiazem of claim 14 to the patient in the evening for effective treatment of the patient's hypertension and/or angina the next morning.

33. A method of treatment of a patient's hypertension and/or angina comprising the administration of the preparation of Diltiazem of claim 15 to the patient in the evening for effective treatment of the patient's hypertension and/or angina the next morning.

34. A method of treatment of a patient's hypertension and/or angina comprising the administration of the preparation of Diltiazem of claim 16 to the patient in the evening for effective treatment of the patient's hypertension and/or angina the next morning.

35. A method of treatment of a patient's hypertension and/or angina comprising the administration of the preparation of Diltiazem of claim 17 to the patient in the evening for effective treatment of the patient's hypertension and/or angina the next morning.

36. A method of treatment of a patient's hypertension and/or angina comprising the administration of the preparation of Diltiazem of claim 18 to the patient in the evening for effective treatment of the patient's hypertension and/or angina the next morning.

37. The preparation of claim 1, 2, 3, 4, 7, 8, 9, 11, 12, 13 or 14 wherein the preparation contains 120 mg of Diltiazem.

38. The preparation of claim 1, 2, 3, 4, 7, 8, 9, 11, 12, 13 or 14 wherein the preparation contains 180 mg of Diltiazem.

39. The preparation of claim 1, 2, 3, 4, 7, 8, 9, 11, 12, 13, or 14 wherein the preparation contains 240 mg of Diltiazem.

40. The preparation of claim 1, 2, 3, 4, 7, 8, 9, 11, 12, 13, or 14 wherein the preparation contains 300 mg of Diltiazem.

41. The preparation of claim 1, 2, 3, 4, 7, 8, 9, 11, 12, 13, or 14 wherein the preparation contains 360 mg of Diltiazem.

42. The preparation of claim 1, 2, 3, 4, 7, 8, 9, 11, 12, 13, or 14 wherein the preparation contains 420 mg of Diltiazem.

43. A method of treatment of a patient's hypertension and/or angina comprising the administration of the preparation of Diltiazem of claim 37, 38, 39, 40, 41 or 42 to the patient in the evening for effective treatment of the patient's hypertension and/or angina the next morning.

44. The preparation of claim 9, 10, 11, 12, 13, 14, 15 or 16 wherein the wetting agent is selected from:

sugars;

saccharose, mannitol, sorbitol;

lecithins;

C₁₂ to C₂₀ fatty acid esters of saccharose, commercialized under the name of sucroesters (Gattefosse, France) or under the name of crodesters (Croda, U.K.) such as sucrose stearate marketed under the trade name of Crodesta;

xylose esters or xylites;

polyoxyethylenic glycerides;

esters of fatty acids and polyoxyethylene (Brijs, Renex and Eumulgines, Henkel, RFA);

sorbitan fatty acid esters (Span, Atlas, U.S.A.);

polyglycides-glycerides and polyglycides-alcohols esters (Gelucires, Gattefosse, France)

Metal salts such as NaCl or sodium lauryl sulphate.

45. The preparation of claim 9 wherein the wetting agent is in association with the diltiazem in the bead and not mixed therewith, the membrane comprises a water-soluble or water dispersible polymer or copolymer such as hydroxypropylmethylcellulose and a water-, acid- and base-insoluble polymer which is a neutral copolymer of acrylic acid ethyl ester and acrylic acid methyl ester such as Eudragit NE30D enabling the bead to be hydrated by the introduction of intestinal fluids into the core hydrating the core and therefore mixing the diltiazem and the wetting agent.

46. A method of treatment of a patient's hypertension and/or angina comprising the administration of the preparation of Diltiazem of claim 44 to the patient in the evening for effective treatment of the patient's hypertension and/or angina the next morning.

47. A method of treatment of a patient's hypertension and/or angina comprising the administration of the preparation of Diltiazem of claim 45 to the patient in the evening for effective treatment of the patient's hypertension and/or angina the next morning.

48. The preparation of claim 9, 10, 11, 12, 13, 14, 15, 16, 44 or 45 in which the core and membrane comprise:

	% W/W
(a) Diltiazem hydrochloride	69 - 73
(b) Microcrystalline cellulose (Avicel ph101)	8 - 9.5
(c) Povidone K30	1 - 2
(d) Sucrose stearate (crodesta F150)	7 - 8
(e) Magnesium stearate NF	0.5 - 2.5
(f) Talc USP	0.5 - 5.0
(g) Titanium dioxide (USP)	0.15 - 0.3
(h) Hydroxypropylmethylcellulose 2910	0.3 - 0.6
(i) Polysorbate 80 (tween)	0.01 - 0.025
(j) Simeticone C emulsion USP (dry of 30%)	0.01 - 0.015
(k) Eudragit NE30 D (dry of 30%)	7 - 11
Purified water USP	0 (used for mixing)

49. A method of treatment of a patient's hypertension and/or angina comprising the administration of the preparation of Diltiazem of claim 46 to the patient in the evening for effective treatment of the patient's hypertension and/or angina the next morning.

50. The preparation of claim 9, 10, 11, 12, 13, 14, 15, 16, 44 or 45 in which the core and membrane comprise:

(i) in the core,

(a) between about 50% and about 85% (% w/w of the total preparation) of Diltiazem or pharmaceutically acceptable salt thereof; and

(b) between about 2% and about 25% wetting agent (% w/w of the total preparation);

together with suitable adjuvants; and

(ii) in the membrane,

(c) between about 0.1% and about 2% of the total preparation of water-soluble and/or water-dispersible polymer such as hydroxypropylmethylcellulose; and

(d) between about 5% and about 20% (% w/w of the preparation) of a neutral copolymer of acrylic acid ethyl ester and acrylic acid methyl ester (such as Eudragit NE30D), together with suitable adjuvants.

51. A method of treatment of a patient's hypertension and/or angina comprising the administration of the preparation of Diltiazem of claim 50 to the patient in the evening for effective treatment of the patient's hypertension and/or angina the next morning.

(i) in the core,

(b) between about 7% and about 8% wetting agent (% w/w of the total preparation);

together with suitable adjuvants; and

(ii) in the membrane,

(c) between about 0.3% and about 0.6% of the total preparation of water-soluble and/or water-dispersible polymer such as hydroxypropylmethylcellulose; and

(d) between about 7% and about 11% (% w/w of the preparation) of a neutral copolymer of acrylic acid ethyl ester and acrylic acid methyl ester (such as Eudragit NE30D), together with suitable adjuvants.

53. A method of treatment of a patient's hypertension and/or angina comprising the administration of the preparation of Diltiazem of claim 52 to the patient in the evening for effective treatment of the patient's hypertension and/or angina the next morning.

54. The preparation of claim 9, 10, 11, 12, 13, 14, 15, 16, 44 or 45 wherein the preparation is a tablet and the tablet comprises microgranules in association with wax placebo beads which wax placebo beads serve to absorb the shock placed on the microgranules of Diltiazem during the tablet process, together with suitable excipients and adjuvants.

55. A method of treatment of a patient's hypertension and/or angina comprising the administration of the preparation of Diltiazem of claim 54 to the patient in the evening for effective treatment of the patient's hypertension and/or angina the next morning.

56. A controlled-release Galenical preparation of pharmaceutically acceptable Diltiazem including the pharmaceutically acceptable salts thereof, suitable for evening dosing every 24 hours containing from about 120 mg to about 540 mg or more of the form of Diltiazem with excipients to provide controlled (sustained) release of the form of Diltiazem from the preparation for providing a C_{max} of Diltiazem in the blood at between about 10 hours and about 15 hours (T_{max}) after administration of the preparation, the preparation being in a sustained-release dosage form in which the Diltiazem is adapted to be control released after administration of the preparation over a period of time wherein the preparation comprises a plurality of microgranules, each microgranule comprising a central core

containing the form of diltiazem coated with a microporous membrane and the central core comprises Diltiazem or pharmaceutically acceptable salt thereof associated with a wetting agent in which the core and membrane comprise:

(i) in the core,

(a) between about 50% and about 85% (% w/w of the total preparation) of Diltiazem or pharmaceutically acceptable salt thereof; and

(b) between about 2% and about 25% wetting agent (% w/w of the total preparation);

together with suitable adjuvants; and

(ii) in the membrane,

(c) between about 0.1% and about 2% of the total preparation of water-soluble and/or water-dispersible polymer such as hydroxypropylmethylcellulose; and

(d) between about 5% and about 20% (% w/w of the preparation) of a neutral copolymer of acrylic acid ethyl ester and acrylic acid methyl ester (such as Eudragit NE30D), together with suitable adjuvants.

57. The preparation of claim 56 wherein the microgranules are in capsule form.
58. The preparation of claim 56 wherein the microgranules are in tablet form.
59. The preparation of claim 56, 57 or 58 wherein the core and membrane comprise:

(i) in the core,

(a) between about 69% and about 73% (% w/w of the total preparation) of Diltiazem or pharmaceutically acceptable salt thereof; and

(b) between about 7% and about 8% wetting agent (% w/w of the total preparation);

together with suitable adjuvants; and

(ii) in the membrane,

(c) between about 0.3% and about 0.6% of the total preparation of water-soluble and/or water-dispersible polymer such as hydroxypropylmethylcellulose; and

(d) between about 7% and about 11% (% w/w of the preparation) of a neutral copolymer of acrylic acid ethyl ester and acrylic acid

methyl ester (such as Eudragit NE30D), together with suitable adjuvants.

60. The preparation of claim 56, 57 or 58 wherein the core and membrane comprise:

	% W/W
(a) Diltiazem hydrochloride	69 - 73
(b) Microcrystalline cellulose (Avicel ph101)	8 - 9.5
(c) Povidone K30	1 - 2
(d) Sucrose stearate (crodesta F150)	7 - 8
(e) Magnesium stearate NF	0.5 - 2.5
(f) Talc USP	0.5 - 5.0
(g) Titanium dioxide (USP)	0.15 - 0.3
(h) Hydroxypropylmethylcellulose 2910	0.3 - 0.6
(i) Polysorbate 80 (tween)	0.01 - 0.025
(j) Simeticone C emulsion USP (dry of 30%)	0.01 - 0.015
(k) Eudragit NE30 D (dry of 30%)	7 - 11
Purified water USP	0 (used for mixing).

61. The preparation of claim 56, 58, 59 or 60 wherein the preparation is a tablet and the tablet comprises microgranules in association with wax placebo beads which wax placebo beads serve to absorb the shock placed on the microgranules of Diltiazem during the tablet process, together with suitable excipients and adjuvants.

[illegible]

ABSTRACT

A controlled-release Galenical preparation of pharmaceutically acceptable Diltiazem including the pharmaceutically acceptable salts thereof, suitable for evening dosing every 24 hours containing from about 120 mg to about 540 mg or more of the form of Diltiazem with excipients to provide controlled (sustained) release of the form of Diltiazem from the preparation for providing a C_{max} of Diltiazem in the blood at between about 10 hours and about 15 hours (T_{max}) after administration of the preparation, the preparation being in a sustained-release dosage form in which the Diltiazem is adapted to be control released after administration of the preparation over a period of time and being adapted to release the Diltiazem


(i) into an aqueous medium at the following rates measured using the method of United States Pharmacopoeia No. XXIII at 100 rpm in 900 ml of water:

- (a) between about 1% and about 15% after 2 hours;
- (b) between about 7% and about 35% after 4 hours;
- (c) between about 30% and about 58% after 8 hours;
- (d) between about 55% and about 80% after 14 hours; and
- (e) and in excess of about 75% after 24 hours.

and/or (ii) into a buffered medium having a pH between about 5.5 and about 6.5, at the following rates measured using the method of United States Pharmacopoeia No. XXIII at 100 rpm in 900ml of the buffered medium:


- (a) between about 1% and about 25% after about 2 hours;
- (b) between about 7% and about 45% after about 4 hours;
- (c) between about 30% and about 68% after about 8 hours;
- (d) in excess of about 75% after about 24 hours.

Figure 1



09465338 . 121799


Figure 2



05463338.121799

Figure 3:

DILTIAZEM DISSOLUTION



09465338 121759

Figure 4




Figure 5




Figure 6


Percent Dissolved [%]

Diltiazem HCl

DISSOLUTION PROFILE

MEDIUM: INTESTINAL FLUID

STRENGTH: 120 mg



09465338 121799

Figure 7





Figure 8



09463333 161799

Declaration, Power Of Attorney And Petition

We, **KENNETH STEPHEN ALBERT** and **PAUL JOSÉ MAES**, declare that we are citizens of the United States of America and Belgium, respectively, residing at **87 Greenwood Lane, Mt. Kisco, N.Y. 10549, United States of America**, and **1242 King's College Drive, Oakville, Ontario, L6M 2T9, Canada**, respectively.

That we verily believe we are the original, first and joint inventors of the subject matter which is claimed and for which a patent is sought on the invention entitled **CHRONOTHERAPEUTIC FORMULATIONS OF DILTIAZEM AND THE ADMINISTRATION THEREOF**, which is described and claimed in the accompanying specification.

That we have reviewed and understand the contents of the above-identified specification, including the Claims as amended by any amendments referred to.

That we do not know and do not believe that this invention was ever known or used in the United States of America before our invention or discovery thereof, or patented or described in any printed publication in any country before our invention or discovery thereof, or more than one year prior to this application.

That this invention was not in public use or on sale in the United States of America for more than one year prior to this application.

That this invention or discovery has not been patented or made the subject of an inventor's certificate issued before the date of this application in any country foreign to the United States of America on an application filed by us or our legal representatives or assigns more than twelve (12) months before this application.

That we acknowledge our duty to disclose information of which we are aware which is material to the examination of this application in accordance with 37 CFR 1.56(a), and that no application for patent or inventor's certificate on this invention or discovery has been filed by us or our representatives or assigns in any country foreign to the United States of America except as follows:

Canadian Patent Application Serial No. _____, filed on December 10, 1999, from which application convention priority is claimed.

That we hereby claim the benefit under Title 35, United States Code, S120 of any United States applications listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in the prior United States application in the manner provided by the first paragraph of Title 35, United States Code, S112, we acknowledge the duty to disclose material information as defined in Title 37, Code of Federal Regulations, S1.56(a) which occurred between the filing date of the prior application and the national or PCT international filing date of this application:

UNITED STATES APPLICATIONS
None

Application Serial No.	Filing Date	Status - Patented/pending/abandoned.
------------------------	-------------	--------------------------------------

And we hereby appoint **IVOR M. HUGHES**, Suite 200, 175 Commerce Valley Drive West, Thornhill, Ontario, L3T 7P6, Canada, Registration Number 27,759; **NEIL H. HUGHES**, Suite 200, 175 Commerce Valley Drive West, Thornhill, Ontario, L3T 7P6, Canada, Registration Number 33,636; and **MARCELO K. SARKIS** Suite 200, 175 Commerce Valley Drive West, Thornhill, Ontario, L3T 7P6, Canada, Registration Number 37,015; as our attorneys or agents to prosecute this application and to transact all business in the Patent Office connected therewith.

Wherefore we pray that Letters Patent be granted to us for the invention or discovery described and claimed in the foregoing specification and claims, and we hereby subscribe our name to the foregoing specification and claims, declaration, power of attorney, and this petition.

We declare further that all statements made herein of our own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issuing thereon.

INVENTOR KENNETH STEPHEN ALBERT
First Name Middle Name Last Name

X Kenneth Stephen Albert
Signature

DATE December 16 1999
Month Day Year

INVENTOR PAUL JOSÉ MAES
First Name Middle Name Last Name

Paul Maes
Signature

DATE December 16 1999
Month Day Year

POST OFFICE ADDRESS 2488 Dunwin Drive
Mississauga, Ontario
Canada
L5L 1J9

654741 88888888